



Tidsskriftet
DEN NORSKE LEGEFORENING

Births in Norwegian midwife-led birth units 2008–10; a population-based study

ORIGINALARTIKKEL

PÅL ØIAN

E-mail: pal.oian@unn.no

Department of Obstetrics and Gynaecology
University Hospital of North Norway
Department of Clinical Medicine
University of Tromsø

Pål Øian (b. 1948) is a specialist in obstetrics and gynaecology, senior consultant and professor. The author has completed the ICMJE form and declares no conflicts of interest.

OLAUG MARGRETE ASKELAND

Division of Health Data and Digitalisation
Norwegian Institute of Public Health
Bergen

Olaug Margrete Askeland (b. 1975) is a statistician (MSc) specialising in data analysis, and a senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

INGER ELISE ENGELUND

Centre for patient reported outcomes data
R&D Department
Bergen Hospital Trust

Inger Elise Engelund (b. 1965) is a sociologist (MA) and a consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

BRIT ROLAND

Department of Hospital Services
The Norwegian Directorate of Health

Brit Roland (b. 1954) is a midwife and senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

MARTA EBBING

Division of Health Data and Digitalisation
Norwegian Institute of Public Health
Bergen

Marta Ebbing (b. 1963) is a specialist in heart diseases and has a PhD in clinical epidemiology. She is medical director of the Health Registries.

The author has completed the ICMJE form and declares no conflicts of interest.

BACKGROUND

Midwife-led birth units form a part of the differentiated and decentralised delivery of maternity care services in Norway. The purpose of this study was to investigate the incidence and characteristics of planned and unplanned births in midwife-led birth units, reasons for transfers to hospital, and the outcomes for mother and child.

MATERIAL AND METHOD

In the period between 2008 and 2010, a supplementary form was used for submitting data to the Medical Birth Registry, in addition to the routine report. These forms were filled in by the attending midwife for 2 514 deliveries in midwife-led birth units, out of a total of 2 556 (98.4%), and for 220 deliveries that had been planned for a midwife-led birth unit, but where the birth took place elsewhere. Data obtained from the completed supplementary forms were then linked with the reports routinely submitted to the Medical Birth Registry, after which the outcomes from births in midwife-led birth units were compared to the outcomes from a low-risk birthing population in hospital.

RESULTS

Among the 2 514 deliveries in midwife-led birth units, 2 320 of them (92.3%) had been planned for this birthplace, while 194 had not (7.7%). Among the planned midwife-led birth units deliveries, the total intrapartum transfer rate was 6.9%. Of these, 19.5% were nulliparous women. In standard midwife-led birth units, 0.4% of all births were operative vaginal deliveries, while in modified midwife-led birth units, the operative delivery rate was 3.5%. Among transfers from a midwife-led birth unit to a hospital, the operative delivery rate was 12.7%. Among children born in a midwife-led birth unit, 0.6% had an Apgar score of < 7 after 5 minutes, compared to 1.0% among children born in the low-risk comparison group in hospital ($p = 0.04$).

INTERPRETATION

Midwife-led birth units should submit outcomes for all women selected for giving birth there, irrespective of whether the delivery actually took place in a midwife-led birth unit or elsewhere.

Maternity care in Norway has been a topic of discussion for decades. Internationally, the trend has been towards centralisation of childbirth services, for both financial and medical reasons. It has been maintained that centralisation ensures better utilisation of financial and staff resources and that the outcomes for mother and child are better, particularly in the case of high-risk deliveries (1, 2). The risk that failings in the maternity care provision may lead to the injury or death of mother or child has been found to be greater in small maternity units (1, 3).

Norway has a scattered population, and the Norwegian parliament, the Storting, has decided that we should provide a differentiated and decentralised system of maternity care. Differentiated delivery of maternity services means that different options must be available, to ensure that low-risk women have an opportunity to give birth in high-tech institutions, midwife-led birth units or at home. In Norway, there has been a three-tier system of maternity care, with institutions categorised on the basis of delivery rates, staffing requirements and risk status (4–6): midwife-led birth units (a minimum of 40 deliveries per year, on-call midwifery staff), hospital maternity wards (a minimum of 400–500 deliveries per year, on-call obstetrician and anaesthetist, midwifery and theatre staff as required) and maternity clinics (a minimum of 1 500 deliveries per year, obstetrician and anaesthetist present, on-call paediatrician, midwifery and theatre staff as required, paediatric ward with an intensive care unit). So-called modified midwife-led birth units (with facilities for performing emergency caesarean sections if transport is considered unsafe) have also been

set up in some places (6).

The Medical Birth Registry routinely receives data about all childbirths from the actual place of delivery, not from the mother's intended place of delivery at the time she went into labour. Consequently, we have only limited knowledge of the incidence of, reasons for, and outcomes of intrapartum and postpartum transfers from midwife-led birth units to hospital.

The study's objective was to clarify whether births in midwife-led birth units were intended to take place there, and how often women who started giving birth in a midwife-led birth unit had to be transferred to a hospital. We also wanted to survey the reasons for hospital transfers, ascertain at what point in the birthing process transfers took place, the means of transport used and the outcomes for these deliveries (sphincter injury, blood loss, Apgar score and survival for children up to five years of age). This information is important to facilitate an assessment of the quality of the selection process for midwife-led birth units and the birthing process there.

Material and method

The material consists of detailed data about births that took place in midwife-led birth units between 1 January 2008 and 31 December 2010. The material also includes data about births for which a midwife-led birth unit had been the intended place of delivery but where the woman was transferred, either intrapartum or postpartum, to a different maternity institution. Data were collected by means of a supplementary form drawn up by the Norwegian Directorate for Health and Social Affairs in partnership with the Norwegian Institute of Public Health in 2007 (fig 1). This was distributed to all maternity institutions in the country (midwife-led birth units, hospital maternity wards and maternity clinics). All district midwives and midwives involved with home births were also informed and the form was sent to them. The form was also to be completed when the outcome of a planned home birth was a hospital transfer and for deliveries in transit that were unrelated to births in midwife-led birth units. Data from the latter events were not included in this study.

Additional form for registering deliveries in midwife-led birth units, deliveries involving transfer from midwife-led birth units, planned home deliveries resulting in a transfer, and deliveries in transit
 + To be completed for all births in midwife-led birth units and for all births that involve a transfer from the home to a midwife-led birth unit, or from a midwife-led birth unit to hospital +

1 Institution number Institution name

2 Mother's full name

3 Address

4 Mother's maiden name (surname) 5 Mother's personal identification no.

6 Where was the intended place of delivery
 At home
 Midwife-led birth unit
 Hospital

7 At home
 Midwife-led birth unit
 Hospital
 In transit

Child's date of birth

If multiple birth, specify no.:

Order, if relevant

8 If delivery took place in transit, was the baby delivered between:
 Home and midwife-led birth unit / hospital
 Midwife-led birth unit and hospital

9 Who was present during the delivery:
 Midwife
 Doctor
 Others, describe:

10 Did the delivery in transit take place in...
 Private car?
 Taxi?
 Ambulance?
 Aeroplane/helicopter?
 Boat?
 Other, describe:

11 If the birth took place at home or in a midwife-led birth unit, but this was not the intended birthplace, indicate the reason (you may tick several options)
 Birth progressed too far
 Strong contractions
 Adverse weather conditions/closed roads
 No transport available
 Other, describe:

12 Transfer from home or midwife-led birth unit prepartum or intrapartum
 Date and time of transfer
 Day Month Year (4 digits) Time

13 Reason for transfer prepartum or intrapartum
 Ruptured membranes and not established labour
 Haemorrhaging
 Exhausted mother/need for pain relief
 Suspected infection
 Abnormal presentation
 Fetal distress/cardiotocography changes
 Slow progress in the opening phase
 Slow progress in the second stage
 Other, describe:

14 If a home birth was planned, for how long (in hours) was the woman in the midwife-led birth unit or at home with midwife present, prior to transfer? hours : minutes

15 Cervical dilation when the transfer decision was made cm

16 For how long (in hours) was the woman in labour prior to transfer? hours : minutes

17 Cervical dilation on departure cm

18 Transfer from home or midwife-led birth unit postpartum
 Date and time of transfer
 Day Month Year (4 digits) Time

19 Reason for transfer postpartum
 Mother Haemorrhaging
 Retained placenta
 Examined at home/ in midwife-led birth unit prior to onward transport to unit/hospital
 Other, describe:

Sphincter rupture
 subtotal/total (grade III)
 incl. ruptured rectal mucous membrane (grade IV)

Child Respiration problems
 Deformities
 Injury/fracture
 Jaundice
 Other, describe:

+ Signature _____ + Date _____ +

Figure 1 Supplementary form used throughout the project period

Data were collected pursuant to the Medical Birth Registry Regulations (7) and the information was managed in accordance with procedures drawn up by the Norwegian Institute of Public Health in its capacity as the registry's data administrator. Data were quality assured and analysed by staff at the Norwegian Institute of Public Health. There is no need to obtain the consent of the registered individuals in order to use information submitted to the Medical Birth Registry, nor is approval by the regional ethics committee (REK) required for analysing this data if the analysis is integral to the administration of the registry.

When the study commenced, there were 14 midwife-led birth units in Norway: Føderiket (the former Rikshospitalet in Oslo), Hallingdal (Ål), Lykkeliten (Rjukan), Tynset and Valdres (Fagernes) administered by the Southern and Eastern Norway Regional Health Authority; Lærdal and Odda governed by the Western Norway Regional Health Authority; and Mosjøen, Brønnøysund, Lofoten, Steigen, Midt-Troms (Finnsnes), Sonjatun (Nordreisa) and Alta governed by the Northern Norway Regional Health Authority. The midwife-led birth units in Lærdal, Odda and Lofoten were modified units throughout the study period. One midwife-led birth unit (Lykkeliten) was closed down in the period.

The supplementary form was completed by the midwife in charge of the delivery at the midwife-led birth unit, even if the birthing woman was transferred to hospital and delivery took place while in transit to or in hospital. Completed forms were submitted to the Norwegian Institute of Public Health where they were scanned and checked. Based on information about the identity of the woman and the date of birth of the child, the data were linked with the relevant standard birth report submitted to the Medical Birth Registry. Missing forms were chased up with the appropriate midwife-led birth unit by telephone.

A total of 27 supplementary forms that were completed in the course of the study period

could not be linked to any routine data submitted to the Medical Birth Registry, mainly due to insufficient information about the identity of women with foreign names. Furthermore, there was not always a perfect match between the option ticked on the supplementary form and the registered birthplace category recorded in the Medical Birth Registry. For the purpose of analysis, we have used the information provided in the supplementary form as the gold standard.

Information from standard and modified midwife-led birth units is provided in three categories:

- births in midwife-led birth units that were intended to take place there
- births in midwife-led birth units that were intended to take place elsewhere
- births that were planned for midwife-led birth units but that took place elsewhere

Furthermore, we show the results for a comparison group consisting of low-risk deliveries in hospitals run by the regional health authorities of Southern and Eastern Norway, Western Norway and Northern Norway during the study period. We have defined low-risk births as all deliveries of infants born after 36 weeks gestation, except deliveries by women with chronic diseases such as hypertension, kidney disease, rheumatoid arthritis or heart disease, or who had suffered pregnancy complications such as gestational diabetes and hypertensive complaints. Also excluded were planned caesarean deliveries and deliveries that were induced.

The analyses were conducted using the software SPSS version 22.0 (IBM Corp., Armonk, NY, US) and R version 3.2.4 (8). In order to investigate whether group differences were statistically significant, a non-parametric significance test (chi-squared test) was used. $P < 0.05$ was considered statistically significant.

Results

A total of 3 546 supplementary forms were received, of which 2 734 referred to childbirth in midwife-led birth units and were thus included in the study (fig 2). Among them, 2 514 forms referred to deliveries that took place in a midwife-led birth unit while 220 forms referred to births for which a midwife-led birth unit had been the intended place of delivery, but which in fact took place elsewhere. Based on the standard reports submitted routinely to the Medical Birth Registry, a total of 2 556 births were reported to have taken place in a midwife-led birth unit, which amounted to 1.4 % of all births recorded in the Medical Birth Registry (9). The coverage of the supplementary forms for deliveries in midwife-led birth units was therefore 98.4 %




Figure 2 Categories of collected data and deliveries included

Table 1 provides an overview of the deliveries included in the study by regional health authority and midwife-led birth unit, as well as the intended and actual place of delivery. The midwife-led birth unit had been the intended place of delivery for 2 320 of the 2 514 deliveries that took place there (92.3%), while 194 (7.7%) had been planned for elsewhere.

Table 1

Births by actual and intended place of delivery, 2008–10.

Midwife-led birth unit, Regional Health Authority	Intended place of delivery: Number (%)		Total ¹
	Midwife-led birth unit	Hospital	
Føderiket (Oslo), Southern and Eastern	344 (100)	0 (0)	344
Hallingdal (Ål), Southern and Eastern	1 (33.3)	2 (66.7)	3
Lykkeliten (Rjukan), Southern and Eastern	21 (77.8)	6 (22.2)	27
Tynset, Southern and Eastern	230 (93.9)	14 (5.7)	245
Valdres, Southern and Eastern	96 (88.1)	13 (11.9)	109
Lærdal, Western	218 (90.1)	22 (9.1)	242
Odda, Western	183 (98.4)	3 (1.6)	186
Alta, Northern Norway	249 (87.4)	36 (12.6)	285
Brønnøysund, Northern Norway	114 (96.6)	4 (3.4)	118
Lofoten, Northern Norway	346 (93.8)	18 (4.9)	369
Midt-Troms (Finnsnes), Northern Norway	308 (91.7)	27 (8.0)	336
Mosjøen, Northern Norway	140 (84.3)	26 (15.7)	166
Sonjatun (Nordreisa), Northern Norway	59 (81.9)	13 (18.1)	72
Steigen, Northern Norway	11 (91.7)	1 (8.3)	12
Deliveries in midwife-led birth units	2 320 (92.3)	185 (7.4)	2 514
Deliveries elsewhere	220 (100)	0 (0.0)	220
Deliveries included in the study	2 540 (92.9)	185 (6.8)	2 734

¹Information about the intended place of delivery was missing for eight deliveries; Lærdal: 2,

Lofoten: 5 and Mid-Troms: 1. One delivery at Tynset was planned as a home birth. These nine births have been included in the total number of deliveries in midwife-led birth units.

Among the 2 540 women who had planned to give birth in a midwife-led birth unit, a total of 174 (6.9 %) were transferred to hospital intrapartum. Among the 2 514 women who gave birth in a midwife-led birth unit, 95 (3.8 %) were transferred to hospital postpartum. Among nulliparous women 19.5 % were transferred intrapartum and 3.8 % postpartum, while among multiparous women the intrapartum transfer rate was 3.4 % and the postpartum transfer rate 3.7 % (data not shown).

Figure 2 shows the actual place of delivery for the 220 births that had been planned for a midwife-led birth unit but which took place elsewhere. There were 26 deliveries in transit, of which 16 took place between the home and the midwife-led birth unit; four in a private car and the rest in an ambulance (road vehicle/plane/helicopter). A midwife was present at 18 of the deliveries in transit, while the ambulance personnel assisted in eight such cases and the father of the child in three such cases. No doctor was reported to be present at any of the deliveries in transit.

Among the reasons listed for unplanned births at home or in a midwife-led birth unit (n = 203) the most common was that the birth had progressed too far for transport to be considered practical (139 cases, 68.5 %), and/or frequent and heavy maternal contractions (71 cases, 35.0 %). In four cases adverse weather conditions /closed roads were cited as the reason and in seven cases there was no suitable means of transport available (data not shown).

Table 2 provides an overview of the most important reasons why women are transferred to hospital before or during a birth that was planned to take place in a midwife-led birth unit. Slow progress in the opening phase was listed as a contributing or sole reason in 67 of 174 such cases (38.5 %).

Table 2

Reasons for prepartum and intrapartum transfers in cases of planned midwife-led birth unit deliveries, 2008–10

Reasons for prepartum and intrapartum transfers, n = 174 Number ¹	
Ruptured membranes without established birth (>24 h)	26
Haemorrhage	0
Exhausted mother /need for pain relief	35
Suspected infection	4
Abnormal position / presentation	16
Fetal distress/ cardiotocography changes	41
Slow progress during the opening phase	67
Slow progress during the second stage	11
Other	16

¹Multiple reasons were listed for some deliveries.

The most important reasons for postpartum transfers included severe maternal perineal tears, haemorrhaging or retained placenta, whereas the most frequent cause in the case of the child was respiration problems (data not shown).

When intrapartum transport to hospital was necessary, it took an average of 10.1 hours (median 8.5 hours, range of variation 0–40 hours) from arrival at the midwife-led birth unit until a transfer decision was made, and the dilation of the cervix was 5 cm (average and median, range of variation 0–10 cm).

Table 3 shows the birthing women's characteristics as well as results from standard and modified midwife-led birth units in the three categories of delivery covered by the study.

Additionally, the same data is specified for a low-risk birthing population in hospital (maternity ward or clinic). There were 7 (0.4 %) operative vaginal deliveries at standard midwife-led birth units, 28 (3.5 %) at modified midwife-led birth units and 28 (12.7 %) among those who were transferred to hospital intrapartum. In the low-risk birthing population in hospital, vacuum or forceps was used in 10.4 % of deliveries, which is a considerably higher rate than in the midwife-led birth units ($p < 0.001$). The caesarean section rate for modified midwife-led birth units was 9.5 %, 11.8 % among those transferred from a midwife-led birth unit to a hospital, and 6.2 % in the comparison group. In the midwife-led birth unit population, 0.9 % of women haemorrhaged more than 1 500 ml, while the rate was 1.5 % in the comparison group ($p = 0.02$). Among the children born at a midwife-led birth unit, 0.6 % had an Apgar score of < 7 after five minutes, compared to 1.0 % of the children in the comparison group born in hospital ($p = 0.04$).

Table 3a

Characteristics, complications and outcomes for planned and unplanned births at midwife-led birth units, for births planned for midwife-led birth units that took place elsewhere and for low-risk births in hospital, 2008–10. Data obtained from supplementary forms linked with data from the Medical Birth Registry (MFR) a) Information on deliveries and b) Information on the condition of neonates.

a		Births in midwife-led birth units, planned				Births in midwife-led birth units, unplanned				Births planned for a midwife-led birth unit, took place elsewhere				Births in hospital, low risk ¹	
Deliveries		2 298 of 2 320 linked with MFR data				189 of 194 linked with MFR data				220 of 220 linked with MFR data				105 358 from MFR	
Woman and pregnancy	Categories	Ordinary (n= 1 557)		Modified (n = 741)		Ordinary (n=141)		Modified (n=48)		Ordinary (n=178)		Modified (n=42)		N	%
		N	%	N	%	N	%	N	%	N	%	N	%		
Parity	0	225	14.5	216	29.1	46	32.6	13	27.1	90	50.6	24	57.1	45	43.4
	1+	1 332	85.5	525	70.9	95	67.4	35	72.9	88	49.4	18	42.9	59	56.6
Gestation	Gestation < 36 weeks	2	0.1	10	1.3	5	3.5	4	8.3	1	0.6	0	0.0	0	0.0
	≥ 36 weeks	1 552	99.7	728	98.2	136	96.5	44	91.7	174	97.8	41	97.6	104	99.1
	Missing	3	0.2	3	0.4	0	0.0	0	0.0	3	1.7	1	2.4	962	0.9
Presentation	Normal, vertex	1 470	94.4	710	95.8	130	92.2	46	95.8	150	84.3	39	92.9	98	93.2
	Breech	19	1.2	14	1.9	2	1.4	0	0.0	6	3.4	0	0.0	1	1.8
	Other	67	4.3	17	2.3	8	5.7	2	4.2	22	12.4	3	7.1	5	5.0
	Missing	1	0.1	0	0.0	1	0.7	0	0.0	0	0.0	0	0.0	18	0.0
Operative deliveries:															
Vacuum	Yes	7	0.4	24	3.2	0	0.0	2	4.2	25	14.0	0	0.0	9	8.7
	Forceps	0	0.0	2	0.3	0	0.0	0	0.0	1	0.6	2	4.8	1	1.7
Caesarean sections:	Elective	0	0.0	44	5.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Acute	1	0.1	27	3.6	0	0.0	3	6.3	21	11.8	5	11.9	6	6.2
	Unspecified	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	8	0.0
Complications:															
Sphincter injury, grade 3/4	Yes	13	0.8	18	2.4	2	1.4	0	0.0	2	1.1	0	0.0	2	2.3
Haemorrhage > 500 ml	500–1 500 ml	58	3.7	48	6.5	12	8.5	2	4.2	15	8.4	4	9.5	12	11.7
	> 1 500 ml, transfusion	10	0.6	9	1.2	3	2.1	1	2.1	5	2.8	0	0.0	1	1.5
	> 500 ml, no further details	15	1.0	0	0.0	0	0.0	0	0.0	5	2.8	0	0.0	264	0.3

¹Low-risk single deliveries after ≥ 36 weeks gestation according to routine reports submitted to the Medical Birth Registry (MFR) for the Regional Health Authorities of Southern and Eastern Norway, Western Norway and Northern Norway for 2008–10. Deliveries by women with chronic illnesses or pregnancy complications were excluded, as were induced

deliveries and planned caesarean sections.

Table 3b

Characteristics, complications and outcomes for planned and unplanned births at midwife-led birth units, for births planned for midwife-led birth units that took place elsewhere and for low-risk births in hospital, 2008–10. Data obtained from supplementary forms linked with data from the Medical Birth Registry (MFR) a) Information on deliveries and b) Information on the condition of neonates.

b		Births in midwife-led birth units, planned				Births in midwife-led birth units, unplanned				Births planned for a midwife-led birth unit, took place elsewhere				Births in hospital, low risk ¹	
Newborn infants ¹		2 298 of 2 320 linked with MFR data				190 of 195 linked with MFR data				220 of 220 linked with MFR data				105 358 from MFR	
Categories		Ordinary (n=1 557)		Modified (n=741)		Ordinary (n=142)		Modified (n=48)		Ordinary (n=178)		Modified (n=42)			
The child		N	%	N	%	N	%	N	%	N	%	N	%	N	%
Weight at birth	< 2 500 g	3	0.2	11	1.5	7	4.9	1	2.1	0	0.0	0	0.0	953	0.9
	2 500–4 500 g	1 515	97.3	701	94.6	130	91.5	40	83.3	172	96.6	40	95.2	101 569	96.4
	> 4 500 g	39	2.5	29	3.9	4	2.8	7	14.6	6	3.4	2	4.8	2 789	2.6
	Missing	0	0.0	0	0.0	1	0.7	0	0.0	0	0.0	0	0.0	47	0.0
Apgar score after 5 min	< 7	1	0.1	10	1.3	1	0.7	2	4.2	2	1.1	1	2.4	1 029	1.0
	≥ 7	1 552	99.7	730	98.5	139	97.9	45	93.8	174	97.8	41	97.6	104 289	99.0
	Missing	4	0.3	1	0.1	2	1.4	1	2.1	2	1.1	0	0.0	40	0.0

¹The number of newborns is higher than the number of births, due to an unplanned delivery of twins in a midwife-led birth unit.

Among the entire midwife-led birth unit population there were three still births, all of which were intrauterine deaths found on arrival. Five live infants died between 29 days and one year of their birth. None of these deaths could be related to complications at birth. None of the children in the midwife-led birth unit population were registered as dead between the ages of one and five.

Discussion

Our findings indicate that risk selection and outcomes for midwife-led birth units are satisfactory. A total of 6.9 % of the women who had planned to give birth in a midwife-led birth unit were transferred intrapartum and 3.8 % were transferred postpartum, due to circumstances relating to the mother or the child. Most transfers were uneventful, and the reasons most commonly cited were slow progress in the opening phase, the need for further pain relief, no delivery imminent more than 24 hours after ruptured membranes, discoloured amniotic fluid or changes to the fetal heart sound. Among nulliparous women who had planned to give birth in a midwife-led unit, 19.5 % were transferred intrapartum. This is a considerable proportion, and nulliparous women who wish to give birth in a midwife-led birth unit must be informed of this. However, the transfer rate in our study is clearly lower than the reported rate for midwife-led birth units in England, where observations indicate that 30–43 % of nulliparous women in midwife-led birth units are transferred to a hospital (10, 11).

The most common reasons for postpartum transfers were retained placenta, suspected tears that required repair by a doctor, or respiratory problems in the newborn infant. The use of vacuum/forceps, the incidence of sphincter injuries and the incidence of Apgar score < 7

after 5 minutes demonstrated satisfactory outcomes for deliveries in midwife-led birth units as well as for maternal transfers.

The reasons why 7.7% of deliveries in midwife-led birth units were intended to take place elsewhere, may be that the women waited too long before setting off to hospital, or that the birth progressed so fast that they would be unable to get to hospital and that the only option available if an unplanned home birth or delivery in transit was to be avoided, was to travel to the midwife-led birth unit. Adverse weather conditions or non-availability of suitable means of transport were rare reasons for unintended deliveries in midwife-led birth units. It is also somewhat surprising that 1.4% of births in midwife-led birth units are breech deliveries.

The organisation of maternity services and the number of maternity institutions in Norway will depend on factors such as maternal accessibility, quality of service, economy and political decision-making. It has been decided that we should offer a differentiated and decentralised service provision in a three-tier system of maternity institutions (midwife-led birth units, hospital maternity wards, maternity clinics) (4–6). These categories were formerly based on delivery rates and staffing requirements, but they are now based on quality requirements (12).

It is important that all types of maternity institutions are familiar with their own results. Information reported to the Medical Birth Registry is recorded under the actual birthplace. Consequently, the data that are routinely submitted to the Medical Birth Registry do not always provide an adequate basis for assessing the quality of midwife-led birth units and small maternity wards that requisition maternal transfers to larger maternity institutions. Each institution's results must be analysed according to the intention-to-treat principle, i.e. outcomes must be reported for all birthing women admitted, including those who are transferred intrapartum or postpartum. If a midwife-led birth unit can demonstrate good results for the deliveries that take place within the institution, but poor results for deliveries in transit, then the quality of the services provided by that institution is not necessarily good. This study sought to examine this particular issue.

The findings show satisfactory results for mother and child following deliveries in midwife-led birth units. Only 0.6% of the children had an Apgar score of < 7 after 5 minutes, and we found no data to suggest that serious injuries or deaths in mother or child were related to the midwife-led birth unit delivery per se. The operative vaginal delivery rate was 0.4% in standard midwife-led birth units. These results match earlier Norwegian and international studies (13–20). The study demonstrated a conspicuously high incidence of caesarean deliveries among planned births in modified midwife-led birth units; contrary to the intention, these were not limited to emergency caesarean sections.

The study sheds light on the reasons for intrapartum and postpartum transfers to hospital, at what point in the birthing process transfers took place, and the means of transport. A good ambulance service is clearly important for the midwife-led birth units.

In 2008, at the start of the project period, there were 55 maternity institutions in Norway, of which 14 were midwife-led birth units. In the course of the study period 1.4% of all childbirths took place in midwife-led birth units. There are currently 47 maternity institutions, of which six are midwife-led birth units. The following eight units have been closed down:

Lykkeliten (2008), Føderiket, Lærdal and Steigen (2011), Valdres and Odda (2013), as well as Hallingdal and Mosjøen (2016). In 2016 a total of 0.7% of all childbirths took place in midwife-led birth units (9). Norwegian health authorities have now recommended the winding up of modified midwife-led birth units (12), and there is currently only one such modified unit in the country (Lofoten).

Some maternity services are provided by institutions similar to midwife-led birth units (low-risk units) at Oslo University Hospital (the ABC clinic, Ullevål hospital), Stavanger University Hospital (Fødeloftet) and Haukeland University Hospital (Storken). These units

were not included in the study, but it will be important for them to be able to document results for birthing women who are transferred from these low-risk units to ordinary maternity wards.

In Denmark, Sweden and Finland, maternity care services are more centralised than in Norway. In these countries there are very few maternity wards with less than 1 000 deliveries per year, and Norway has more than twice as many maternity institutions relative to the birth rate. Geography and settlement patterns may well be different, but this can hardly explain the large differences in the number of maternity institutions.

It is a strength of our study that it is population-based, national in scope and prospective. We were able to supplement the collected material with data routinely submitted to the Medical Birth Registry. The supplementary forms provided details about virtually all deliveries in midwife-led birth units during the study period, including the reasons for transfers to hospital, at what point in the birthing process transfers took place, what means of transport was used, and information about deliveries in transit. Furthermore, the supplementary forms provided detailed information about births that had been planned for a midwife-led birth unit, but that took place elsewhere.

An important weakness of the study is that the number of deliveries included is low, and that all findings associated with rare events such as severe maternal haemorrhage, still births, very low Apgar scores and neonatal death, must be interpreted with care.

Conclusion

The study's results suggest that the risk selection and outcomes for midwife-led birth units are satisfactory. While only few multiparous women who planned to give birth in a midwife-led birth unit had to be transferred intrapartum to hospital, nearly one in five nulliparous women were transferred. Most of these transfers were uneventful. The study provided knowledge about delivery outcomes that are required for the quality assessment of selection for and delivery in midwife-led birth units. All low-risk maternity units and small maternity wards should be in a position to document such knowledge.

MAIN MESSAGE

In the period between 2008 and 2010, 1.4 % of all childbirths in Norway took place in midwife-led birth units.

Among the women who gave birth in midwife-led birth units and had planned to do so, 6.9 % were transferred to a hospital intrapartum while 3.8 % were transferred postpartum.

Among births in standard midwife-led birth units, 0.4 % were operative vaginal deliveries, while the operative delivery rate was 3.5 % in modified midwife-led birth units.

Among the children born in midwife led birth units, 0.6 % had an Apgar score of < 7 after 5 minutes, compared to 1.0 % among the children born in a low-risk population in hospital.

REFERENCES:

1. Milland M, Christoffersen JK, Hedegaard M. The size of the labor wards: is bigger better when it comes to patient safety? *Acta Obstet Gynecol Scand* 2013; 92: 1271 - 6. [PubMed][CrossRef]
2. Dietl J. Trends in birth statistics in Europe. *Z Geburtshilfe Neonatol* 2002; 206: 48 - 50. [PubMed][CrossRef]
3. Johansen LT, Øian P. Barn som dør eller får alvorlig skade under fødsel. *Tidsskr Nor Legeforen* 2011; 131: 465 - 8. [CrossRef]
4. Faglige krav til fødeinstitusjoner. Statens helsetilsyn. Utredningsserie 1-97. <https://helsedirektoratet.no/Lists/Publikasjoner/Attachments/245/Et-trygt-fodetilbud-kvalitetskrav-til->

fodselsomsorgen-IS-1877.pdf (26.9.2017).

5. St. meld. nr. 43 (1999-2000). Om akuttmedisinsk beredskap.

<https://www.regjeringen.no/no/dokumenter/stmeldnr-43-1999-2000/id193493/sec1> (26.9.2017).

6. St. meld. nr. 12 (2008-2009) En gledelig begivenhet.

<https://www.regjeringen.no/no/dokumenter/stmeld-nr-12-2008-2009/id545600/> (26.9.2017).

7. FOR-2001-12-21-1483. Forskrift om innsamling og behandling av helseopplysninger i Medisinsk fødselsregister (Medisinsk fødselsregisterforskriften).

<https://lovdata.no/dokument/SF/forskrift/2001-12-21-1483> (13.3.2018).

8. The R Project for Statistical Computing The R Project for Statistical Computing. 2018. The R Project for Statistical Computing The R Project for Statistical Computing. <http://www.R-project.org> (13.3.2018).

9. Medisinsk fødselsregister (MFR) og Abortregisteret. <http://statistikkbank.fhi.no/mfr/> (13.3.2018).

10. Brocklehurst P, Hardy P, Hollowell J et al. Perinatal and maternal outcomes by planned place of birth for healthy women with low risk pregnancies: the Birthplace in England national prospective cohort study. *BMJ* 2011; 343: d7400. [PubMed][CrossRef]

11. Hundley VA, Cruickshank FM, Lang GD et al. Midwife managed delivery unit: a randomised controlled comparison with consultant led care. *BMJ* 1994; 309: 1400 - 4. [PubMed][CrossRef]

12. Et trygt fødetilbud. Kvalitetskrav til fødselsomsorgen. Helsedirektoratet 12/2010.

<https://helsedirektoratet.no/Lists/Publikasjoner/Attachments/245/Et-trygt-fodetilbud-kvalitetskrav-til-fodselsomsorgen-IS-1877.pdf> (26.9.2017).

13. Schmidt N, Abelsen B, Øian P. Deliveries in maternity homes in Norway: results from a 2-year prospective study. *Acta Obstet Gynecol Scand* 2002; 81: 731 - 7. [PubMed][CrossRef]

14. Holt J, Vold IN, Backe B et al. Child births in a modified midwife managed unit: selection and transfer according to intended place of delivery. *Acta Obstet Gynecol Scand* 2001; 80: 206 - 12. [PubMed]

15. Lukasse M, Øian P, Aamodt G. En jordmorledet fødeenhet. *Tidsskr Nor Laegeforen* 2006; 126: 170 - 2. [PubMed]

16. Bernitz S, Rolland R, Blix E et al. Is the operative delivery rate in low-risk women dependent on the level of birth care? A randomised controlled trial. *BJOG* 2011; 118: 1357 - 64. [PubMed][CrossRef]

17. Hollowell J, Li Y, Bunch K et al. A comparison of intrapartum interventions and adverse outcomes by parity in planned freestanding midwifery unit and alongside midwifery unit births: secondary analysis of 'low risk' births in the birthplace in England cohort. *BMC Pregnancy Childbirth* 2017; 17: 95. [PubMed][CrossRef]

18. Christensen LF, Overgaard C. Are freestanding midwifery units a safe alternative to obstetric units for low-risk, primiparous childbirth? An analysis of effect differences by parity in a matched cohort study. *BMC Pregnancy Childbirth* 2017; 17: 14. [PubMed][CrossRef]

19. National Institute for Health and Care Excellence. Intrapartum care for healthy women and babies. NICE clinical guideline 190, 2014.

<https://www.nice.org.uk/guidance/cg190/resources/intrapartum-care-for-healthy-women-and-babies/pdf-35109866447557> (26.9.2017).

20. Delgado Nunes V, Gholitabar M, Sims JM et al. Intrapartum care of healthy women and their babies: summary of updated NICE guidance. *BMJ* 2014; 349: g6886. [PubMed][CrossRef]

Published: 12 June 2018. *Tidsskr Nor Legeforen*. DOI: 10.4045/tidsskr.17.0842

Received 28.9.2017, first revision submitted 23.12.2017, accepted 13.3.2018.

© The Journal of the Norwegian Medical Association 2020. Downloaded from tidsskriftet.no